

Test Name	Laboratory	Specimen Type	Test Schedule	Reference Range	Effective Date	Comments
-----------	------------	---------------	---------------	-----------------	----------------	----------

1,25 diOH Vitamin D (see 1,25-Dihydroxy Vitamin D, Serum/Plasma)

--	--	--	--	--	--	--

Test Name	Laboratory	Specimen Type	Test Schedule	Reference Range	Effective Date	Comments
1,25-Dihydroxy Vitamin D, Serum/Plasma Calcitriol 1,25 diOH Vitamin D	Endocrinology	<p>Adult: 5 mL Gold top Vacutainer tube</p> <p>Pediatric: 0-2 years: 2 x 0.5 mL Red or Gold top Microtainers 2-10 years: 3 mL Red top Vacutainer tube</p> <p>Light Green top tubes (Li- heparin) or Lavender top tubes (EDTA) are also acceptable GENERAL LABORATORY REQUISITION</p>	Monday-Friday 0800-1600	60-208 pmol/L	2015-01-05	<p>The test for 1,25 di-OH vitamin D is <u>not</u> useful to assess sufficiency of vitamin D from diet, supplements, and endogenous synthesis. The 25-OH vitamin D test would be more appropriate for that purpose.</p> <p><u>Appropriate Indications for 1,25 di-OH Vitamin D Testing:</u></p> <p>RENAL FAILURE</p> <p>RENAL TUBULOPATHY</p> <p>SARCOIDOSIS</p> <p>UNEXPLAINED HIGH PARATHYROID HORMONE LEVEL</p> <p>PEDIATRIC ENDOCRINOLOGY CONCERN (E.g. vitamin D-dep (more...))</p>

Test Name	Laboratory	Specimen Type	Test Schedule	Reference Range	Effective Date	Comments
1. Erythrocyte Total Panel (Total, Essential and Toxic) 2. Essential Erythrocyte Panel 3. Toxic Erythrocyte Panel 4. Total Panel	Trace Elements	Reference # 368381-BD Royal Blue K2-EDTA Vacutainer TRACE ELEMENTS REQUISITION	Batched analysis	Find individual Reference Ranges here:		Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here:
11-Deoxycortisol,Serum/Plasma	Core (all campuses)	6 mL Red top Vacutainer tube or 4.5 mL Lavender top tube Pediatric: 0-2 years: 0.5 mL Red Microtainer 2-10 years: 2 mL Red top GENERAL LABORATORY REQUISITION	As required	See report for therapeutic ranges	2008-12-16	Referred out Monday - Thursday Hospitals In-Common Laboratory Inc.
14C Breath Test (see <u>H. Pylori Breath Test</u> test only available to Grey Bruce, Owen Sound and St. Mary's, Kitchener)						
17 Alpha Hydroxy Progesterone (see <u>17-Hydroxyprogesterone, Serum</u>)						
17 Beta Estradiol (see <u>Estradiol, Plasma/Serum</u>)						
17-Hydroxy Progesterone (see <u>17-Hydroxyprogesterone, Serum</u>)						

Test Name	Laboratory	Specimen Type	Test Schedule	Reference Range	Effective Date	Comments
17-Hydroxyprogesterone, Serum 17-OH Progesterone 17-Hydroxy Progesterone 17 Alpha Hydroxy Progesterone	Core	<p>Adult: 6 mL Red top Vacutainer tube</p> <p>Pediatric: 0-2 years: 2 x 0.5 mL Red top Microtainers 2-10 years: 3 mL Red top Vacutainer tube</p> <p>Avoid blood collection tubes with separator gels (Gold top Vacutainer tubes)</p> <p>Light Green (Li-Heparin) or Lavender (EDTA) top tubes are NOT acceptable GENERAL LABORATORY REQUISITION</p>	Referred out Monday - Friday	<p><14 days: <4.9 nmol/L 14 days - <1 year: <3.5 nmol/L 1 - <12 years: <1.2 nmol/L 12 - <14 years: <2.1 nmol/L 14 - <16 years: <4.3 nmol/L 16 - <19 years: <4.0 nmol/L</p> <p>Male: ≥19 years: <6.0 nmol/L</p> <p>Female: Follicular phase: <5.6 nmol/L Midcycle: <6.8 nmol/L Luteal phase: <8.6 nmol/L Post-menopausal: <1.4 nmol/L</p>	2017-11-07	
17-OH Progesterone (see 17-Hydroxyprogesterone, Serum)						