

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|----------------------|--|---|-----------------|----------------|---|
| Q fever Serology Coxiella burnetti Serology | Microbiology (VH) | 5 mL Gold top or 6 mL Red top Vacutainer tube PUBLIC HEALTH LABORATORY TEST REQUISITION | Referred out weekdays to the Public Health Laboratory. | | 2010-09-13 | Q Fever has an incubation period of approximately 2 to 3 weeks. Positive antibody results should be correlated with clinical findings. |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|----------------------|--|------------------------|------------------|----------------|---|
| QF-PCR Rapid Aneuploidy Detection | Cytogenetics (VH) | Amniotic Fluid CVS Fetal blood / Cord Blood Skin Fetal Tissue Products of Conception CYTOGENETI CS REQUISITION (must include patient's name, address, Ontario Health Insurance Number, PIN (if applicable), originating location, Dr's full name and address, and pertinent clinical information. | Batched as required | See final report | 2016-03-22 | The Cytogenetics Lab is staffed from 0700- 1700 (Monday-Friday), Ext. 78974 (office),or 77022(lab) or 75714(Sr. Tech) For additional information please refer to the Molecular Diagnostics Laboratory N/A See final report |
| Quantitative Cytomegalovirus (see <u>CMV PCR</u>) | | | | | | |
| Quantitative Epstein Barr Virus (see <u>EBV PCR</u>) | | | | | | |
| Quantitative RNA (see <u>Hepatitis C RNA - Quantitative</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---------------------|------------|--|-------------------------------------|-----------------|----------------|----------|
| Quinidine Biquin | Core | 6 mL Red top Vacutainer tube Avoid gel- separator tubes GENERAL LABORATORY REQUISITION | Referred out Monday- Thursday | 6.0-15.0 µmol/L | | |