

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|----------------------------------|--|--|---|----------------|---|
| T-Cell Lymphoma TCRG Gene clonality assay | Molecular Diagnostics | EDTA blood/bone marrow MOLECULAR DIAGNOSTIC REQUISITION | As required, Monday to Friday 0800-1600h | See report | | |
| T. vulgaris (see <u>Farmers Lung IgG Antibodies, Serum</u>) | | | | | | |
| Tacrolimus FK506 Prograf | Toxicology/Speci al Chemistry | 4 mL K ₂ or K ₃ EDTA Lavender top Vacutainer tube Pediatric: 0-2 yrs: 0.5 mL Lavender 2-10 yrs: 3 mL Lavender top GENERAL LABORATORY REQUISITION | Samples are tested Monday- Saturday. Specimens received in the Core lab after 10:00am will be processed the next working day. | No established reference range. Concentrations are measured in ng/mL. | 2009-02-27 | Call the Toxicology/Special Chemistry Laboratory for more information at: (519) 685-8500 x 64664 option 3. Thoroughly mix the blood sample after procurement. |
| Taenia solium Serology (see <u>Cysticercosis Serology</u>) | | | | | | |
| Tau Protein (see <u>Beta-2 Transferrin</u>) | | | | | | |
| Tay-Sachs Disease (see <u>Beta-N-Acetylhexosaminidase %A, A, A+B, Leukocyte/Plasma/Fibroblasts</u>) | | | | | | |
| TB Culture (see <u>Mycobacterium Culture</u>) | | | | | | |
| TBG (see <u>Thyroxine Binding Globulin, Serum</u>) | | | | | | |
| TBII (see <u>Anti-Thyroid Stimulating Hormone Receptor Antibodies, Serum</u>) | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|--|---------------------------------------|-----------------|----------------|----------|
| TCOM (see <u>Complement Total, Serum</u>) | | | | | | |
| TCRG Gene clonality assay (see <u>T-Cell Lymphoma</u>) | | | | | | |
| Tegretol (see <u>Carbamazepine, Serum/Plasma-Total</u>) | | | | | | |
| Telo peptide-N (see <u>N-Telo peptide, Urine</u> no longer available- see C-Telo peptide) | | | | | | |
| Tempra (see <u>Acetaminophen</u>) | | | | | | |
| Teriflunomide (see <u>Leflunomide</u>) | | | | | | |
| Testicular Biopsy | Pathology | Tissue biopsy Powerchart: eOrder choosing appropriate specimen. See Identification of Clinical Specimens | Refer to Collection Information | See report | 2019-04-09 | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|------------------------------------|------------|---|---------------|--|----------------|---|
| Testosterone (Total), Plasma/Serum | Core | <p>Adult: 4.5 mL Light Green top (Li-Heparin) Vacutainer tube</p> <p>Pediatric: 0-2 years: 0.5 mL Light Green top (Li-Heparin) Microtainer 2-10 years: 3 mL Light Green top (Li-Heparin) Vacutainer tube</p> <p>Red, Gold, or Lavender (EDTA) top tubes are also acceptable</p> <p>GENERAL LABORATORY REQUISITION</p> | As required | <p><u>Male:</u> Tanner stage 1: ≤ 0.4 nmol/L Tanner stage 2: ≤ 15.0 nmol/L Tanner stage 3: 2.3 - 27.0 nmol/L Tanner stage 4: 6.2 - 26.5 nmol/L Tanner stage 5: 6.5 - 30.6 nmol/L 20 - 49 years: 8.6 - 29.0 nmol/L ≥ 50 years: 6.7 - 25.7 nmol/L</p> <p><u>Female:</u> Tanner stage 1: ≤ 0.4 nmol/L Tanner stage 2: ≤ 0.4 nmol/L Tanner stage 3: ≤ 0.8 nmol/L Tanner stage 4: ≤ 0.9 nmol/L Tanner stage 5: ≤ 1.3 nmol/L 20 - 49 years: ≤ 1.7 nmol/L ≥ 50 years: ≤ 1.4 (more...)</p> | 2018-03-06 | <p>Nandrolone (19-nortestosterone) interferes with this assay.</p> <p>Biotin may interfere with this test. Samples should not be taken from patients receiving high biotin doses (i.e. > 5 mg/day) until at least 8 hours after the last biotin administration.</p> <p>In isolated cases, elevated testosterone levels can be seen in samples from female patients with end stage renal disease (ESRD).</p> |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|----------------|--|--|--|----------------|---|
| Tetanus Serology | Microbiology | 5 mL Gold top Vacutainer tube PUBLIC HEALTH LABORATORY TEST REQUISITION | Referred out weekdays to Public Health Laboratory. | | 2017-10-02 | Testing is available through the Public Health Laboratory only for the rare event of an adverse reaction to the Tetanus vaccine or the possibility of humoral immunodeficiency. |
| Tg (see <u>Thyroglobulin, Serum/Plasma</u>) | | | | | | |
| Thallium, Erythrocytes | Trace Elements | Reference number 368381 - HMMS# 11073 - 6mL K2-EDTA Royal Blue Vacutainer tube TRACE ELEMENTS REQUISITION | Batched analysis | SI Units (Reported on Patient Chart): 0.00-0.13 nmol/L Conventional Units: 0.000-0.026 µg/L | | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|----------------|----------------|---|------------------|---|----------------|--|
| Thallium,Urine | Trace Elements | 24 hour urine collected in an unused 24 hour urine container or random urine TRACE ELEMENTS REQUISITION | Batched analysis | <u>SI Units:</u> Random Urine: 0-1.96 nmol/L nmol/mol creatinineAgeFe maleMale0-110- 2330-22212-190- 1420-13820-290- 1600-12230-390- 1940-14340-490- 2300-15150-590- 2720-17360-690- 2680-18170-790- 2800-196≥800- 3430-222 24 Hour Urine: 0-2.94 nmol/d <u>Conventional Units:</u> Random Urine: 0-0.400 µg/L ng/g creatinineAgeFe maleMale0-110- 4210-40 (more...) | | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|----------------|--|------------------------------------|--|----------------|---|
| Thallium, Whole blood | Trace Elements | Reference number 368381 - HMMS# 11073 - 6mL K2-EDTA Royal Blue Vacutainer tube TRACE ELEMENTS REQUISITION | Batched analysis | SI Units (Reported on Patient Chart): 0.00-0.19 nmol/L Conventional Units: 0.000-0.039 µg/L | | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |
| Theo-Dur (see <u>Theophylline, Serum</u>) | | | | | | |
| Theophylline, Serum Aminophylline Theo-Dur Somophyllin Elixophyllin Xanthine | Core | Adult: 6 mL Red top Vacutainer tube Pediatric: 0-2 years: 2 x 0.5 mL Red top Microtainers 2-10 years: 3 mL Red top Vacutainer tube GENERAL LABORATORY REQUISITION | Referred out Monday Thursday | 55 - 110 mol/L | 2010-08-31 | Bronchodilator drug used in treatment of reversible bronchoconstriction. Aminophylline is a salt of theophylline (80-85% by weight of this salt is theophylline). Toxic: >110 mol/L Critical Value to be phoned to nurse or physician immediately |
| Thermoactinomyces vulgaris (see <u>Farmers Lung IgG Antibodies, Serum</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|------------|--|------------------------------|---|----------------|----------|
| Thiamine Diphosphate (see <u>Vitamin B1, Whole Blood</u>) | | | | | | |
| Thiamine Pyrophosphate (Whole Blood) (see <u>Vitamin B1, Whole Blood</u>) | | | | | | |
| Thiocyanate, Serum/Plasma Sulfocyanate Nitroprusside | Core | 6 mL Red top Vacutainer tube or 4 mL Lavender EDTA Vacutainer tube Avoid gel-separator tubes GENERAL LABORATORY REQUISITION | Referred out Monday-Thursday | Therapeutic: Less than 0.5 mmol/L Toxic: Greater than 1.7 mmol/L | 1998-09-01 | |
| Thiopurine Methyltransferase: Genotype TPMT Genotype | Core | 4 mL Lavender EDTA Vacutainer tube GENERAL LABORATORY REQUISITION | Referred out Monday-Thursday | Results interpreted on report | 2015-08-25 | |
| Thiopurine Methyltransferase: Phenotype TPMT Phenotype | Core | 4 mL Lavender EDTA Vacutainer tube GENERAL LABORATORY REQUISITION | Referred out Monday-Thursday | Low Activity: < 10 nmol/Hbh Intermediate: 10 - 40 nmol/Hbh Normal Activity: > 40 nmol/Hbh | 2015-08-25 | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|-------------------|---|------------------------------|--|----------------|---|
| Thioridazine and Metabolite, Serum/Plasma Mellaril | Core | 5 mL Gold top Vacutainer tube Avoid gel-separator tubes GENERAL LABORATORY REQUISITION | Referred out Monday-Thursday | Thioridazine steady-state serum concentration during chronic oral administration of 400 mg daily: 140 - 2600 ng/mL Therapeutic steady-state concentrations may overlap levels associated with toxicity. Mesoridazine therapeutic range: 100 - 1400 ng/mL | | Thioridazine is a phenothiazine for treating psychosis. Monitoring its level is important to optimize therapy, to avoid toxicity, and to assure compliance. Toxic: Greater than 27 µmol/L |
| Throat Culture | Microbiology (VH) | Throat, tonsil MICROBIOLOGY REQUISITION | Daily | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|--------------------------|---|---|-----------------|----------------|---|
| Thrombophilia (associated with Factor V deficiency) Factor V Familial Thrombophilia Prothrombin/Factor V Prothrombin | Molecular Diagnostics | 1 x 4 mL K ₂ or K ₃ EDTA Lavender top Vacutainer tube-may be a frozen aliquot. MOLECULAR DIAGNOSTIC REQUISITION | As Required Monday - Friday 0800 - 1600 h | See report | | For more information click on: Molecular Diagnostic Laboratory N/A Resistance to activated protein C (APC) is a major cause of familial thrombophilia, and can be corrected by an anticoagulant activity expressed by purified factor V. It has been suggested that a point mutation in the gene coding for factor V is responsible for APC resistance (PMID:7909098, 8208267). This point mutation, (F5:c.1601G>A), occurring towards the 3' end of exon 10 of the factor V gene, is predicted to cause a missense mutation in the APC cleavage site, (F5:p.Arg534Gln) and confers an inc (more...) |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|--------------------------|--|---|-----------------|----------------|---|
| Thrombophilia (Prothrombin) Prothrombin Gene | Molecular Diagnostics | Whole blood-1 x 4 mL Lavender EDTA top Vacutainer tube MOLECULAR DIAGNOSTIC REQUISITION | As required Monday - Friday 0800 - 1600 h | See report | | For more information click on: Molecular Diagnostic Laboratory N/A At least five genetic defects,(PMID:9299960) accounting for approximately 15% of families with inherited thrombophilia, have been established as risk factors for venous thrombosis. There are protein C, protein S, and antithrombin deficiencies, and represent defects in the anticoagulant pathways of blood coagulation. Two other genetic risk factors, resistance to activated protein C associated with the factor V Leiden mutation and increased prothrombin associated with the prothrombin 20210 A allele (more...) |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|---|---------------|-----------------|----------------|---|
| Thrombophilia Screen (see <u>Hypercoagulable Screen</u>) | | | | | | |
| Thyrocalcitonin (see <u>Calcitonin, Fine Needle Aspirate, Calcitonin, Serum</u>) | | | | | | |
| Thyroglobulin, Fine Needle Aspirate | Core | <p>Two samples: The first is uncontaminated Plasma-Lyte and serves as a blank.</p> <p>The second is 1 mL of Plasma-Lyte that has been used to rinse the biopsy needle as described below.</p> <p>GENERAL LABORATORY REQUISITION</p> | As required | ≤ 1.0 g/L | 2017-07-04 | <p>The limit of quantitation of the assay is 0.1 g/L. If a blank result of > 0.1 g/L were to be obtained, a technical investigation would be performed prior to reporting of result.</p> <p>The purpose of the blank fluid is to confirm that the fluid being used to rinse the needle does not have a matrix effect in the assay.</p> <p>An elevated thyroglobulin level in the fine needle aspirate may indicate the presence of metastatic differentiated thyroid cancer.</p> |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--------------------------------|------------|---|---------------|---|----------------|--|
| Thyroglobulin, Serum/Plasma Tg | Core | <p>Adult: 5 mL Gold top Vacutainer tube</p> <p>Pediatric: 0-2 years: 0.5 mL Red or Gold top Microtainer 2-10 years: 3 mL Red top Vacutainer tube</p> <p>Lavender top (EDTA) tubes are also acceptable</p> <p>Light Green top (Li-Heparin) tubes are NOT acceptable</p> <p>GENERAL LABORATORY REQUISITION</p> | As required | <p>Thyroidectomized patients: < 0.2 g/L</p> <p>Normal population: 3.5 - 77.0 g/L</p> | 2017-07-04 | <p>Biotin may interfere with this test. Samples should not be taken from patients receiving high biotin doses (i.e. > 5 mg/day) until at least 8 hours after the last biotin administration.</p> <p>In the follow-up of thyroidectomized patients, a concentration > 0.2 g/L is suspicious for local or metastatic disease.</p> <p>In non-thyroidectomized patients, a concentration greater than the reference interval is not diagnostic of cancer.</p> <p>This assay is to be used to monitor patients who have had thyroidectomy for cancer. It is not recommended (more...)</p> |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|---------------|---------------|-----------------|----------------|----------|
| Thyroid (see <u>Image-Guided Fine Needle Aspirate Cytology, Non-Image Guided Fine Needle Aspiration Biopsy for Cytology</u>) | | | | | | |
| Thyroid Antibodies (see <u>Anti-Thyroglobulin Antibodies, Serum/Plasma</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|------------|---|---------------|---|----------------|---|
| Thyroid Stimulating Hormone, Plasma/Serum TSH TSH-3 Thyrotropin | Core | <p>Adult: 4.5 mL Green top Vacutainer tube</p> <p>Pediatric: 0-2 years: 0.5 mL Light Green top (Li-Heparin) Microtainer 2-10 years: 3 mL Light Green top (Li-Heparin) Vacutainer tube</p> <p>Red, Gold, or Lavender (EDTA) top tubes are also acceptable GENERAL LABORATORY REQUISITION</p> | As required | <p>0 6 days: 0.70 15.2 mIU/L</p> <p>6 days 3 months: 0.72 11.0 mIU/L</p> <p>3 12 months: 0.73 8.35 mIU/L</p> <p>1 6 years: 0.70 5.97 mIU/L</p> <p>6 11 years: 0.60 4.84 mIU/L</p> <p>11 20 years: 0.51 4.30 mIU/L</p> <p>> 20 years: 0.27 4.20 mIU/L</p> | 2008-11-15 | <p>TSH should be the initial test to screen for clinically-suspected hypothyroidism or hyperthyroidism. If TSH is below the lower cut-off, FT4 and FT3 testing will be performed reflexively by the laboratory. If TSH is between the lower and upper cut-offs, no FT4 or FT3 testing will be performed reflexively. If TSH is above the upper cut-off, FT4 testing will be performed reflexively by the laboratory. These cut-offs are the TSH reference intervals in children and the optimal cut-offs to predict abnormal FT4 levels in adults.</p> <p>The TSH cut-offs are: 2 <6 years: <0.70 or >5.97 mIU/L (more...)</p> |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|------------|--|--------------------------------|--|----------------|----------|
| Thyroid Stimulating Immunoglobulin (see <u>Anti-Thyroid Stimulating Hormone Receptor Antibodies, Serum</u>) | | | | | | |
| Thyrotropin (see <u>Thyroid Stimulating Hormone, Plasma/Serum</u>) | | | | | | |
| Thyrotropin Binding Inhibitor Immunoglobulin (see <u>Anti-Thyroid Stimulating Hormone Receptor Antibodies, Serum</u>) | | | | | | |
| Thyrotropin Receptor Antibodies (see <u>Anti-Thyroid Stimulating Hormone Receptor Antibodies, Serum</u>) | | | | | | |
| Thyroxine Binding Globulin, Serum TBG | Core | 5 mL Gold top Vacutainer GENERAL LABORATORY REQUISITION | Referred out Monday - Thursday | 0-11 months: 315-685 nmol/L 1 year - 9 years: 278-500 nmol/L 10 years and over: 260-575 nmol/L | | |
| TIBC (see <u>Unsaturated Iron Binding Capacity</u>) | | | | | | |
| Ticks (see <u>Ova and Parasites-Ticks/Arthropods</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|------------|----------------|---|------------------|--|----------------|--|
| Tin, Urine | Trace Elements | 24 hour urine collected in an unused 24 hour urine container or random urine TRACE ELEMENTS REQUISITION | Batched analysis | <u>SI Units:</u> Random Urine: 4.2-42.2 nmol/L µmol/mol creatinineAgeFe maleMale0-110.5-5.00.5-4.812-190.3-3.10.3-3.020-290.3-3.50.3-2.630-390.4-4.20.3-3.140-490.5-5.00.3-3.250-590.6-5.90.4-3.760-690.6-5.80.4-3.970-790.6-6.00.4-4.2≥800.7-7.40.5-4.8 24 Hour Urine: 8.4-59.0 nmol/d <u>Conventional Units:</u> Random Urine: 0.5-5.0 µg/L (more...) | | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|---------------------|--|---|-----------------|----------------|----------|
| Tinea (see <u>Fungus Culture-Dermatophytes</u>) | | | | | | |
| Tip Culture (see <u>Vascular Tip Culture</u>) | | | | | | |
| Tissue / Fluid Investigation | Flow Cytometry (VH) | Lymph Node Biopsy Fine Needle Aspirate CSF Body Fluids Referred-In Samples: FLOW CYTOMETRY REQUISITION | Monday-Friday 0800-1600 Fridays or prior to STAT holidays, routine specimens must be received in the Flow Cytometry Laboratory at Victoria Hospital by 1300. For after hours and weekend requests, page the Hematologist on-call at (519) 685-8500 x 14999. | | 2014-04-21 | |
| Tissue Culture | Microbiology (VH) | Tissue, biopsy MICROBIOLOGY REQUISITION | Daily | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|-----------------|---|----------------------|---|----------------|--|
| Tissue for Immunofluorescence (IF) - Skin, Conjunctival biopsy, Buccal mucosa | Immunopathology | Tissue biopsy PowerChart: E-order choosing appropriate specimen. See Identification of Clinical Specimens. | Tuesday and Thursday | Negative | | |
| Tissue Transglutaminase Antibody (see Anti-Tissue Transglutaminase Antibodies (IgA), Serum/Plasma) | | | | | | |
| Titanium, Erythrocytes | Trace Elements | Reference number 368381 - HMMS# 11073 - 6mL K2-EDTA Royal Blue Vacutainer tube TRACE ELEMENTS REQUISITION | Batched analysis | SI Units (Reported on Patient Chart): 0.0-23.0 nmol/L Conventional Units: 0.0-1.1 µg/L | | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |
| Titanium, Plasma | Trace Elements | Reference number 368381 - HMMS# 11073 - 6mL K2-EDTA Royal Blue Vacutainer tube TRACE ELEMENTS REQUISITION | Batched analysis | SI Units (Reported on Patient Chart): 0.00-7.9 nmol/L Conventional Units: 0.00-0.38 µg/L | | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|----------------|--|------------------|---|----------------|--|
| Titanium, Serum | Trace Elements | Reference number 368380 - HMMS# 260 - 6 mL Non Additive Royal Blue Vacutainer tube TRACE ELEMENTS REQUISITION | Batched analysis | SI Units (Reported on Patient Chart): 0.0-5.8 nmol/L Conventional Units: 0.00-0.28 µg/L | | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |
| Titanium, Whole Blood | Trace Elements | Reference number 368381 - HMMS# 11073 - 6mL K2-EDTA Royal Blue Vacutainer tube TRACE ELEMENTS REQUISITION | Batched analysis | SI Units (Reported on Patient Chart): 0.0-20.9 nmol/L. Conventional Units: 0.0-1.0 µg/L | 2008-09-19 | Reference Ranges are based on Non-Occupationally exposed population. Find Interpretive Comment and Clinical Information here: |
| TNT - High sensitivity (see <u>High Sensitivity Troponin T</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|---|---------------|--|----------------|----------|
| Tobramycin, Serum/Plasma Aminoglycosides | Core (VH) | 4.5 mL Green (Lithium Heparin) top Vacutainer tube Pediatric: 0-2 years: 0.5 mL Green Microtainer 2-10 years: 2 mL Green top tube GENERAL LABORATORY REQUISITION | Daily | <u>Therapeutic Range</u> Trough (Pre Dose): < = 1.4 mg/L Peak (Post Dose): 6-10 mg/L | 2010-07-12 | |
| Tofranil (see <u>Imipramine, Serum/Plasma</u>) | | | | | | |
| Topamax (see <u>Topiramate, Serum/Plasma</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|------------------------------|--|------------------------------|-----------------|----------------|----------|
| Topiramate, Serum/Plasma Topamax | Toxicology/Special Chemistry | 6 mL Red or 6 mL Green or 4 mL Lavender top Vacutainer tube Pediatric: 2-10 yrs: 2 mL Red or Dark Green top 0-2 yrs: Red 0.5pk. or 0.6 Green pk. GENERAL LABORATORY REQUISITION | Monday - Friday 0800-1600 | 15-60 µmol/L | 2006-06-01 | |
| Total (conjugated and unconjugated) metanephrines (see <u>Metanephrines, Urine</u>) | | | | | | |
| Total Complement Function Assay (see <u>Complement Total, Serum</u>) | | | | | | |
| Total Homocysteine (see <u>Homocysteine</u>) | | | | | | |
| Total PSA (see <u>Prostate Specific Antigen, Plasma/Serum</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|-------------------|--|---|--|----------------|--|
| Toxocara Serology | Microbiology (VH) | Blood-5 mL Gold or 6 mL Red top Vacutainer tube PUBLIC HEALTH LABORATORY TEST REQUISITION | Referred out weekdays to the Public Health Laboratory in Toronto. | | 2010-09-28 | Adequate clinical and epidemiological information must accompany the specimen. |
| Toxoplasma (see <u>Ova and Parasites-Blood and Tissue</u>) | | | | | | |
| Toxoplasma Serology - Toxoplasma IgG/IgM | Core (UH) | 5 mL Gold or 6 mL Red top Vacutainer tube GENERAL LABORATORY REQUISITION | 24/7 | See report Indeterminate Toxo. IgG results will be forwarded to Public Health for confirmation on weekdays. | 2006-07-01 | |
| TPMT Genotype (see <u>Thiopurine Methyltransferase: Genotype</u>) | | | | | | |
| TPMT Phenotype (see <u>Thiopurine Methyltransferase: Phenotype</u>) | | | | | | |
| TRAb (see <u>Anti-Thyroid Stimulating Hormone Receptor Antibodies, Serum</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|----------------|--|------------------|---|----------------|----------|
| Trace Elements Creatinine, Urine Creatinine, Urine- Random for Trace Elements Creute Creatinine Trace Elements | Trace Elements | 24 hour urine collected in an unused 24 hour urine container or random urine TRACE ELEMENTS REQUISITION | Batched analysis | <u>Creatinine</u> <u>Reference</u> <u>Ranges used for</u> <u>random urine</u> <u>calculations</u> <u>for Trace</u> <u>Elements</u> SI Units mmol/L AgeFemaleRangeMean0-112.0-14.78.412-193.6-25.613.820-291.9-23.112.230-391.2-21.410.140-490.8-18.28.550-591.1-15.37.260-691.2-15.57.370-791.3-15.47.0≥800.9-10.65.7 AgeMaleRangeMean0-113.5-15.08.8 (more...) | 2012-06-25 | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|---|-------------------------|-----------------------------------|----------------|----------|
| Trace Elements Magnesium, Urine (see Magnesium for Trace Elements, Urine) | | | | | | |
| Transferrin Isoelectric Focussing, Serum/Plasma | Core | 5 mL Gold top Vacutainer is preferable or 6 mL Green (Sodium Heparinized) top Vacutainer tube is acceptable GENERAL LABORATORY REQUISITION | Referred out Wednesdays | A descriptive report will be sent | 2006-09-06 | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|---------------------|--|---------------|-----------------|----------------|---|
| Transferrin Saturation UIBC Iron Overload Screen | Core (all campuses) | 4.5 mL Light Green (Li Heparin) Vacutainer tube GENERAL LABORATORY REQUISITION | As required | 11.0-56.0% | | <p>The test is not valid and may be misleading in patients receiving iron therapy.</p> <p>Useful in the investigation of suspected iron deficiency or iron overload. Serum ferritin is the preferred method for assessing iron stores.</p> <p>Assessment of patients with acute iron poisoning.</p> <p>In cases of iron deficiency, the transferrin saturation is lowered; with iron overload, it is increased.</p> |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|-------------------------|------------|--|---------------|-----------------|----------------|---|
| Transferrin, Plasma TRF | Core | <p>Adult: 4.5 mL Green (Lithium Heparin) top Vacutainer</p> <p>Serum from a 5 mL Gold top or 6 mL Red top is also acceptable.</p> <p>Pediatric: 0-2 years: 0.5 mL Light Green top (Li-Heparin) Microtainer 2-10 years: 3 mL Light Green top (Li-Heparin) Vacutainer tube</p> <p>GENERAL LABORATORY REQUISITION</p> | Daily | 2.00-3.60 g/L | 2010-01-11 | <p>Turbidimetric assays not suitable for measurement of highly lipemic or hemolytic samples or samples containing high levels of circulating immune complexes.</p> <p>Transferrin quantitation not to be confused with transferrin saturation. To screen for chronic iron overload diseases, hemochromatosis, please order Transferrin saturation.</p> |

Transfusion ABC (see [HLA Transfusion Typing](#))

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|-------------------|---|------------------------------|------------------|----------------|--|
| Transfusion Reaction Investigation | Blood Transfusion | 6 mL Pink (EDTA) top Vacutainer or Donor Tag (SJHC) BLOOD TRANSFUSION REQUISITION or Electronic order | Daily Urgent, if indicated | See report | | System Code: Determined by Transfusion Reaction Course. Further blood transfusion should not occur until investigation is complete N/A Results will be interpreted by Hematologist or designate. |
| Transglutaminase Antibody (see <u>Anti-Tissue Transglutaminase Antibodies (IgA), Serum/Plasma</u>) | | | | | | |
| Transthyretin (see <u>Prealbumin, Serum</u>) | | | | | | |
| Trazodone Desyrel | Core | 4 mL Dark Green (Lithium Heparin) top Vacutainer tube Avoid gel-separator tubes GENERAL LABORATORY REQUISITION | Referred out Monday-Thursday | 2.2 - 4.3 µmol/L | 2005-07-01 | |
| TRF (see <u>Transferrin, Plasma</u>) | | | | | | |
| Trichinella (see <u>Ova and Parasites-Blood and Tissue</u>) | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|-------------------|--|---|-----------------|----------------|---|
| Trichinella Serology | Microbiology (VH) | Blood-5 mL Gold or 6 mL Red top Vacutainer tube PUBLIC HEALTH LABORATORY TEST REQUISITION | Referred out weekdays to the Public Health Laboratory in Toronto. | | 2010-09-28 | Adequate clinical and epidemiological information must accompany the specimen. Clearly indicate on label and requisition "TRICHINOSIS SEROLOGY". |
| Tricyclic Screen Tricyclics, Plasma | Core | 4.5 mL Green (Lithium Heparin) top Vacutainer tube (preferred) or 5 mL Gold top Vacutainer tube Pediatric: 0-2 years: 0.5 mL Green Microtainer 2-10 years: 2 mL Green top tube GENERAL LABORATORY REQUISITION | As required | Negative | 2009-06-04 | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|--|---------------|--|----------------|--|
| Tricyclics, Plasma (see <u>Tricyclic Screen</u>) | | | | | | |
| Triglycerides, Plasma | Core | 4.5 mL Green top Vacutainer Pediatric: 0-2 years: 0.5 mL Green top Microtainer 2-10 years: 3 mL Green top GENERAL LABORATORY REQUISITION | As required | * See interpretation of lipid profile Increased risk of cardiovascular disease at triglycerides level greater than 2.0 mmol/L; increased risk of acute pancreatitis at triglycerides level greater than 10.0 mmol/L (Lancet 2014;384:626-635). | 2008-11-15 | Cholesterol target levels are dependent upon patient 10-year risk of coronary artery disease (Can J Cardiol 2013;29:151-67). Dicynone (Etamsylate) at therapeutic concentrations may lead to false low results and Ascorbic acid and calcium dobesilate cause artificially low triglyceride results. Intralipid is directly measured as analyte in this assay and leads to high triglyceride results. Triglyceride: Suggested optimal plasma triglyceride concentration is <1.50 mmol/L. Triglyceride level >10.00 mmol/L is a risk factor for pancreatitis |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|----------------------|---|------------------------------|---|----------------|----------|
| Trimipramine, Serum/ Plasma Surmontil | Core | 2 x 6 mL Red top Vacutainer tube or 2 x Lavender top Vacutainer tubes Avoid gel-separator tubes GENERAL LABORATORY REQUISITION | Referred out Monday-Thursday | 170-1000 nmol/L | 2005-07-01 | |
| Tripeptidyl Peptidase 1, Dried Blood Spot/Fibroblast CLN2 Peptidase LINCL Batten Disease | Biochemical Genetics | 1. Blood dot on PKU-type card 2. Fibroblast 1. GENERAL LABORATORY REQUISITION 2. BIOCHEMICAL GENETICS LAB REQUISITION | As required | Dried Blood Spot: 2.6-7.8 nmol/24 hr/3 mm disc Fibroblast: 47-114 nmol/hr/mg protein | 2008-06-10 | |
| Triple Marker (see <u>Maternal Serum Screen (for Open Neural Tube Defect and Down Syndrome Risks)</u>) | | | | | | |
| Triple Screen (see <u>Maternal Serum Screen (for Open Neural Tube Defect and Down Syndrome Risks)</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|--|------------|---------------|---------------|-----------------|----------------|----------|
| Triptil (see <u>Protriptyline, Serum</u> TESTING ON HOLD as of 11/09/15) | | | | | | |
| Tropheryma whipplei (see <u>Whipple's Disease</u>) | | | | | | |
| Troponin T - High sensitivity (see <u>High Sensitivity Troponin T</u>) | | | | | | |
| Trypanosoma (see <u>Ova and Parasites-Blood and Tissue</u>) | | | | | | |
| Trypanosomiasis Screen (see <u>Blood Parasite Screen</u>) | | | | | | |
| Trypsin-like immunoreactivity (see <u>Trypsin/Trypsinogen, Serum</u>) | | | | | | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|---|------------------------------|-------------------|----------------|----------|
| Trypsin/Trypsinogen, Serum Immunoreactive trypsin Immunoreactive trypsinogen Trypsin-like immunoreactivity | Core | <p>Adult: 5 mL Gold Vacutainer tube</p> <p>Pediatric: 0-2 years: 2 x 0.5 mL Red or Gold top Microtainers 2-10 years: 3 mL Red top Vacutainer tube</p> <p>Light Green (Li-Heparin) or Lavender (EDTA) top tubes are NOT acceptable GENERAL LABORATORY REQUISITION</p> | Referred out Monday - Friday | 180.5 885.3 ng/mL | 2018-08-07 | |
| | | | | | | |

| Test Name | Laboratory | Specimen Type | Test Schedule | Reference Range | Effective Date | Comments |
|---|------------|---|------------------------------|-----------------|----------------|----------|
| Tryptase, Serum/Plasma | Core | 5 mL Gold top Vacutainer tube or 6 mL Red top Vacutainer tube Plasma is also acceptable. 4.5 mL Green Vacutainer (Li Heparin) Pediatric: 0-2 years: Red 0.5 mL Microtainer 2-10 years: 2 mL Red top tube GENERAL LABORATORY REQUISITION | Referred out Monday-Thursday | 3.8 11.4 g/L | 2009-02-27 | |
| TSH (see <u>Thyroid Stimulating Hormone, Plasma/Serum</u>) | | | | | | |
| TSH-3 (see <u>Thyroid Stimulating Hormone, Plasma/Serum</u>) | | | | | | |
| TSI (see <u>Anti-Thyroid Stimulating Hormone Receptor Antibodies, Serum</u>) | | | | | | |
| tTGAB (see <u>Anti-Tissue Transglutaminase Antibodies (IgA), Serum/Plasma</u>) | | | | | | |
| TTR (see <u>Familial Amyloidotic Polyneuropathy-TTR</u>) | | | | | | |
| Tularensis antibody (see <u>Francisella tularensis Serology</u>) | | | | | | |